



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C07H 21/04, C07K 14/705, C12N 15/09, 15/63, C12Q 1/68</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 00/11015</b> <b>(43) International Publication Date:</b> <b>2 March 2000 (02.03.00)</b>																								
<b>(21) International Application Number:</b> <b>PCT/US99/19351</b> <b>(22) International Filing Date:</b> <b>24 August 1999 (24.08.99)</b>  <b>(30) Priority Data:</b> <table border="0"> <tr> <td>60/097,638</td> <td>24 August 1998 (24.08.98)</td> <td>US</td> </tr> <tr> <td>60/097,659</td> <td>24 August 1998 (24.08.98)</td> <td>US</td> </tr> <tr> <td>60/099,618</td> <td>9 September 1998 (09.09.98)</td> <td>US</td> </tr> <tr> <td>60/102,092</td> <td>28 September 1998 (28.09.98)</td> <td>US</td> </tr> <tr> <td>60/109,978</td> <td>25 November 1998 (25.11.98)</td> <td>US</td> </tr> <tr> <td>60/113,645</td> <td>23 December 1998 (23.12.98)</td> <td>US</td> </tr> <tr> <td>60/113,646</td> <td>23 December 1998 (23.12.98)</td> <td>US</td> </tr> <tr> <td>09/379,246</td> <td>23 August 1999 (23.08.99)</td> <td>US</td> </tr> </table>		60/097,638	24 August 1998 (24.08.98)	US	60/097,659	24 August 1998 (24.08.98)	US	60/099,618	9 September 1998 (09.09.98)	US	60/102,092	28 September 1998 (28.09.98)	US	60/109,978	25 November 1998 (25.11.98)	US	60/113,645	23 December 1998 (23.12.98)	US	60/113,646	23 December 1998 (23.12.98)	US	09/379,246	23 August 1999 (23.08.99)	US	<b>(74) Agent:</b> SPRUNGER, Suzanne, A.; American Home Products Corporation, Patent & Trademark Dept. - 2B, One Campus Drive, Parsippany, NJ 07054 (US).  <b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
60/097,638	24 August 1998 (24.08.98)	US																								
60/097,659	24 August 1998 (24.08.98)	US																								
60/099,618	9 September 1998 (09.09.98)	US																								
60/102,092	28 September 1998 (28.09.98)	US																								
60/109,978	25 November 1998 (25.11.98)	US																								
60/113,645	23 December 1998 (23.12.98)	US																								
60/113,646	23 December 1998 (23.12.98)	US																								
09/379,246	23 August 1999 (23.08.99)	US																								
<b>(71) Applicant:</b> ALPHAGENE, INC. [US/US]; 260 West Cummings Park, Woburn, MA 01801 (US).  <b>(72) Inventors:</b> VALENZUELA, Dario; 260 West Cummings Park, Woburn, MA 01801 (US). YUAN, Olive; 260 West Cummings Park, Woburn, MA 01801 (US). HOFFMAN, Heidi; 260 West Cummings Park, Woburn, MA 01801 (US). HALL, Jeff; 260 West Cummings Park, Woburn, MA 01801 (US). RAPIEJKO, Peter; 260 West Cummings Park, Woburn, MA 01801 (US).		<b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>																								
<b>(54) Title:</b> SECRETED PROTEINS AND POLYNUCLEOTIDES ENCODING THEM  <b>(57) Abstract</b>  Novel polynucleotides and the proteins encoded thereby are disclosed.																										

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

What is claimed is:

1. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:
  - (a) the nucleotide sequence of SEQ ID NO:1;
  - (b) the nucleotide sequence of SEQ ID NO:1 from nucleotide 683 to nucleotide 934;
  - (c) the nucleotide sequence of the full-length protein coding sequence of clone vb11\_1 deposited with the ATCC under accession number 98846;
  - (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vb11\_1 deposited with the ATCC under accession number 98846;
  - (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:2;
  - (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:2, the fragment comprising eight contiguous amino acids of SEQ ID NO:2;
  - (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
  - (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:1.
2. The polynucleotide of claim 1 wherein said polynucleotide is operably linked to at least one expression control sequence.
3. A host cell transformed with the polynucleotide of claim 2.
4. The host cell of claim 3, wherein said cell is a mammalian cell.
5. A process for producing a protein encoded by the polynucleotide of claim 2, which process comprises:

- (a) the amino acid sequence of SEQ ID NO:2;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:2, the fragment comprising eight contiguous amino acids of SEQ ID NO:2; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vb11\_1 deposited with the ATCC under accession number 98846;
- the protein being substantially free from other mammalian proteins.

10. The protein of claim 9, wherein said protein comprises the amino acid sequence of SEQ ID NO:2.

11. A composition comprising the protein of claim 9 and a pharmaceutically acceptable carrier.

12. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:3;
- (b) the nucleotide sequence of SEQ ID NO:3 from nucleotide 63 to nucleotide 482;
- (c) the nucleotide sequence of SEQ ID NO:3 from nucleotide 201 to nucleotide 482;

(d) the nucleotide sequence of the full-length protein coding sequence of clone vb12\_1 deposited with the ATCC under accession number 98846;

(e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vb12\_1 deposited with the ATCC under accession number 98846;

(f) the nucleotide sequence of a mature protein coding sequence of clone vb12\_1 deposited with the ATCC under accession number 98846;

(g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vb12\_1 deposited with the ATCC under accession number 98846;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:4;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:4, the fragment comprising eight contiguous amino acids of SEQ ID NO:4;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:3.

13. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:4;

(b) a fragment of the amino acid sequence of SEQ ID NO:4, the fragment comprising eight contiguous amino acids of SEQ ID NO:4; and

(c) the amino acid sequence encoded by the cDNA insert of clone vb12\_1 deposited with the ATCC under accession number 98846;

the protein being substantially free from other mammalian proteins.

14. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

of clone vb14\_1 deposited with the ATCC under accession number 98846;

(e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vb14\_1 deposited with the ATCC under accession number 98846;

(f) the nucleotide-sequence-of-a mature protein coding sequence of clone vb14\_1 deposited with the ATCC under accession number 98846;

(g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vb14\_1 deposited with the ATCC under accession number 98846;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:6;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:6, the fragment comprising eight contiguous amino acids of SEQ ID NO:6;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:5.

15. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:6;

(b) a fragment of the amino acid sequence of SEQ ID NO:6, the fragment comprising eight contiguous amino acids of SEQ ID NO:6; and

(c) the amino acid sequence encoded by the cDNA insert of clone vb14\_1 deposited with the ATCC under accession number 98846; the protein being substantially free from other mammalian proteins.

16. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:7;
- (b) the nucleotide sequence of SEQ ID NO:7 from nucleotide 82 to nucleotide 294;
- (c) the nucleotide sequence of SEQ ID NO:7 from nucleotide 109 to nucleotide 294;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone ve11\_1 deposited with the ATCC under accession number 98846;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone ve11\_1 deposited with the ATCC under accession number 98846;
- (f) the nucleotide sequence of a mature protein coding sequence of clone ve11\_1 deposited with the ATCC under accession number 98846;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone ve11\_1 deposited with the ATCC under accession number 98846;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:8;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:8, the fragment comprising eight contiguous amino acids of SEQ ID NO:8;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:7.

- (c) the nucleotide sequence of SEQ ID NO:9 from nucleotide 116 to nucleotide 468;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vf2\_1 deposited with the ATCC under accession number 98846;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vf2\_1 deposited with the ATCC under accession number 98846;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vf2\_1 deposited with the ATCC under accession number 98846;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vf2\_1 deposited with the ATCC under accession number 98846;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:10;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:10, the fragment comprising eight contiguous amino acids of SEQ ID NO:10;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and



(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:9.

19. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:10;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:10, the fragment comprising eight contiguous amino acids of SEQ ID NO:10; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vf2\_1 deposited with the ATCC under accession number 98846;
- the protein being substantially free from other mammalian proteins.

20. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:11;
- (b) the nucleotide sequence of SEQ ID NO:11 from nucleotide 124 to nucleotide 1641;
- (c) the nucleotide sequence of SEQ ID NO:11 from nucleotide 262 to nucleotide 1641;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vg2\_1 deposited with the ATCC under accession number 98846;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vg2\_1 deposited with the ATCC under accession number 98846;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vg2\_1 deposited with the ATCC under accession number 98846;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vg2\_1 deposited with the ATCC under accession number 98846;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:12;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:12, the fragment comprising eight contiguous amino acids of SEQ ID NO:12;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:11.

21. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:12;

(b) a fragment of the amino acid sequence of SEQ ID NO:12, the fragment comprising eight contiguous amino acids of SEQ ID NO:12; and

(c) the amino acid sequence encoded by the cDNA insert of clone vg2\_1 deposited with the ATCC under accession number 98846;

the protein being substantially free from other mammalian proteins.

22. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:13;

(b) the nucleotide sequence of SEQ ID NO:13 from nucleotide 380 to nucleotide 892;

(c) the nucleotide sequence of SEQ ID NO:13 from nucleotide 416 to nucleotide 892;

(d) the nucleotide sequence of the full-length protein coding sequence of clone vj1\_1 deposited with the ATCC under accession number 98846;

(e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vj1\_1 deposited with the ATCC under accession number 98846;

(f) the nucleotide sequence of a mature protein coding sequence of clone vj1\_1 deposited with the ATCC under accession number 98846;

(g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vj1\_1 deposited with the ATCC under accession number 98846;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:14;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:14, the fragment comprising eight contiguous amino acids of SEQ ID NO:14;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:13.

23. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:14;

(b) a fragment of the amino acid sequence of SEQ ID NO:14, the fragment comprising eight contiguous amino acids of SEQ ID NO:14; and

(c) the amino acid sequence encoded by the cDNA insert of clone vj1\_1 deposited with the ATCC under accession number 98846;

the protein being substantially free from other mammalian proteins.

24. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:15;

(b) the nucleotide sequence of SEQ ID NO:15 from nucleotide 62 to nucleotide 1057;

(c) the nucleotide sequence of SEQ ID NO:15 from nucleotide 659 to nucleotide 1057;

(d) the nucleotide sequence of the full-length protein coding sequence of clone v11\_1 deposited with the ATCC under accession number 98846;

- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vl1\_1 deposited with the ATCC under accession number 98846;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vl1\_1 deposited with the ATCC under accession number 98846;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vl1\_1 deposited with the ATCC under accession number 98846;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:16;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:16, the fragment comprising eight contiguous amino acids of SEQ ID NO:16;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:15.

25. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:16;
- (b) a fragment of the amino acid sequence of SEQ ID NO:16, the fragment comprising eight contiguous amino acids of SEQ ID NO:16; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vl1\_1 deposited with the ATCC under accession number 98846;

the protein being substantially free from other mammalian proteins.

26. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:17;

(b) the nucleotide sequence of SEQ ID NO:17 from nucleotide 74 to nucleotide 529;

(c) the nucleotide sequence of SEQ ID NO:17 from nucleotide 140 to nucleotide 529;

(d) the nucleotide sequence of the full-length protein coding sequence of clone vk2\_1 deposited with the ATCC under accession number 98838;

(e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vk2\_1 deposited with the ATCC under accession number 98838;

(f) the nucleotide sequence of a mature protein coding sequence of clone vk2\_1 deposited with the ATCC under accession number 98838;

(g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vk2\_1 deposited with the ATCC under accession number 98838;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:18;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:18, the fragment comprising eight contiguous amino acids of SEQ ID NO:18;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:17.

27. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:18;

(b) a fragment of the amino acid sequence of SEQ ID NO:18, the fragment comprising eight contiguous amino acids of SEQ ID NO:18; and

(c) the amino acid sequence encoded by the cDNA insert of clone vk2\_1 deposited with the ATCC under accession number 98838;

the protein being substantially free from other mammalian proteins.

28. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:19;
- (b) the nucleotide sequence of SEQ ID NO:19 from nucleotide 174 to nucleotide 3170;
- (c) the nucleotide sequence of SEQ ID NO:19 from nucleotide 1098 to nucleotide 3170;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vb21\_1 deposited with the ATCC under accession number 98862;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vb21\_1 deposited with the ATCC under accession number 98862;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vb21\_1 deposited with the ATCC under accession number 98862;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vb21\_1 deposited with the ATCC under accession number 98862;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:20;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:20, the fragment comprising eight contiguous amino acids of SEQ ID NO:20;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:19.

29. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:20;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:20, the fragment comprising eight contiguous amino acids of SEQ ID NO:20; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vb21\_1 deposited with the ATCC under accession number 98862;
- the protein being substantially free from other mammalian proteins.

30. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:21;
- (b) the nucleotide sequence of SEQ ID NO:21 from nucleotide 74 to nucleotide 1453;
- (c) the nucleotide sequence of SEQ ID NO:21 from nucleotide 224 to nucleotide 1453;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vc35\_1 deposited with the ATCC under accession number 98862;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vc35\_1 deposited with the ATCC under accession number 98862;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vc35\_1 deposited with the ATCC under accession number 98862;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vc35\_1 deposited with the ATCC under accession number 98862;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:22;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:22, the fragment comprising eight contiguous amino acids of SEQ ID NO:22;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees

C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:21.

31. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:22;
- (b) a fragment of the amino acid sequence of SEQ ID NO:22, the fragment comprising eight contiguous amino acids of SEQ ID NO:22; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vc35\_1 deposited with the ATCC under accession number 98862;

the protein being substantially free from other mammalian proteins.

32. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:23;
- (b) the nucleotide sequence of SEQ ID NO:23 from nucleotide 135 to nucleotide 368;
- (c) the nucleotide sequence of SEQ ID NO:23 from nucleotide 243 to nucleotide 368;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vc36\_1 deposited with the ATCC under accession number 98862;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vc36\_1 deposited with the ATCC under accession number 98862;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vc36\_1 deposited with the ATCC under accession number 98862;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vc36\_1 deposited with the ATCC under accession number 98862;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:24;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:24, the fragment comprising eight contiguous amino acids of SEQ ID NO:24;



(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:23.

33. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:24;

(b) a fragment of the amino acid sequence of SEQ ID NO:24, the fragment comprising eight contiguous amino acids of SEQ ID NO:24; and

(c) the amino acid sequence encoded by the cDNA insert of clone vc36\_1 deposited with the ATCC under accession number 98862;

the protein being substantially free from other mammalian proteins.

34. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:25;

(b) the nucleotide sequence of SEQ ID NO:25 from nucleotide 370 to nucleotide 1662;

(c) the nucleotide sequence of the full-length protein coding sequence of clone vc38\_1 deposited with the ATCC under accession number 98862;

(d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vc38\_1 deposited with the ATCC under accession number 98862;

(e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:26;

(f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:26, the fragment comprising eight contiguous amino acids of SEQ ID NO:26;

(g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and

(h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:25.

35. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:26;

(b) a fragment of the amino acid sequence of SEQ ID NO:26, the fragment comprising eight contiguous amino acids of SEQ ID NO:26; and

(c) the amino acid sequence encoded by the cDNA insert of clone vc38\_1 deposited with the ATCC under accession number 98862;

the protein being substantially free from other mammalian proteins.

36. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:27;

(b) the nucleotide sequence of SEQ ID NO:27 from nucleotide 105 to nucleotide 365;

(c) the nucleotide sequence of SEQ ID NO:27 from nucleotide 147 to nucleotide 365;

(d) the nucleotide sequence of the full-length protein coding sequence of clone vc39\_1 deposited with the ATCC under accession number 98862;

(e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vc39\_1 deposited with the ATCC under accession number 98862;

(f) the nucleotide sequence of a mature protein coding sequence of clone vc39\_1 deposited with the ATCC under accession number 98862;

(g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vc39\_1 deposited with the ATCC under accession number 98862;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:28;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:28, the fragment comprising eight contiguous amino acids of SEQ ID NO:28;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:27.

37. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:28;

(b) a fragment of the amino acid sequence of SEQ ID NO:28, the fragment comprising eight contiguous amino acids of SEQ ID NO:28; and

(c) the amino acid sequence encoded by the cDNA insert of clone vc39\_1 deposited with the ATCC under accession number 98862;

the protein being substantially free from other mammalian proteins.

38. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:29;

(b) the nucleotide sequence of SEQ ID NO:29 from nucleotide 35 to nucleotide 1066;

(c) the nucleotide sequence of SEQ ID NO:29 from nucleotide 128 to nucleotide 1066;

(d) the nucleotide sequence of the full-length protein coding sequence of clone vc40\_1 deposited with the ATCC under accession number 98862;

(e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vc40\_1 deposited with the ATCC under accession number 98862;

(f) the nucleotide sequence of a mature protein coding sequence of clone vc40\_1 deposited with the ATCC under accession number 98862;

(g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vc40\_1 deposited with the ATCC under accession number 98862;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:30;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:30, the fragment comprising eight contiguous amino acids of SEQ ID NO:30;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:29.

39. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:30;

(b) a fragment of the amino acid sequence of SEQ ID NO:30, the fragment comprising eight contiguous amino acids of SEQ ID NO:30; and

(c) the amino acid sequence encoded by the cDNA insert of clone vc40\_1 deposited with the ATCC under accession number 98862;

the protein being substantially free from other mammalian proteins.

40. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:31;

(b) the nucleotide sequence of SEQ ID NO:31 from nucleotide 38 to nucleotide 553;

(c) the nucleotide sequence of SEQ ID NO:31 from nucleotide 104 to nucleotide 553;

- (d) the nucleotide sequence of the full-length protein coding sequence of clone vc46\_1 deposited with the ATCC under accession number 98862;
  - (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vc46\_1 deposited with the ATCC under accession number 98862;
  - (f) the nucleotide sequence of a mature protein coding sequence of clone vc46\_1 deposited with the ATCC under accession number 98862;
  - (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vc46\_1 deposited with the ATCC under accession number 98862;
  - (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:32;
  - (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:32, the fragment comprising eight contiguous amino acids of SEQ ID NO:32;
  - (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
  - (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:31.
41. A protein comprising an amino acid sequence selected from the group consisting of:
- (a) the amino acid sequence of SEQ ID NO:32;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:32, the fragment comprising eight contiguous amino acids of SEQ ID NO:32; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vc46\_1 deposited with the ATCC under accession number 98862;
- the protein being substantially free from other mammalian proteins.

42. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:33;
- (b) the nucleotide sequence of SEQ ID NO:33 from nucleotide 164 to nucleotide 2548;
- (c) the nucleotide sequence of SEQ ID NO:33 from nucleotide 242 to nucleotide 2548;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vc49\_1 deposited with the ATCC under accession number 98862;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vc49\_1 deposited with the ATCC under accession number 98862;
- (f) the nucleotide sequence of a mature protein-coding sequence of clone vc49\_1 deposited with the ATCC under accession number 98862;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vc49\_1 deposited with the ATCC under accession number 98862;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:34;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:34, the fragment comprising eight contiguous amino acids of SEQ ID NO:34;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:33.

43. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:34;
- (b) a fragment of the amino acid sequence of SEQ ID NO:34, the fragment comprising eight contiguous amino acids of SEQ ID NO:34; and

(c) the amino acid sequence encoded by the cDNA insert of clone vc49\_1 deposited with the ATCC under accession number 98862; the protein being substantially free from other mammalian proteins.

44. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:35;
- (b) the nucleotide sequence of SEQ ID NO:35 from nucleotide 150 to nucleotide 776;
- (c) the nucleotide sequence of SEQ ID NO:35 from nucleotide 246 to nucleotide 776;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vc50\_1 deposited with the ATCC under accession number 98862;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vc50\_1 deposited with the ATCC under accession number 98862;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vc50\_1 deposited with the ATCC under accession number 98862;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vc50\_1 deposited with the ATCC under accession number 98862;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:36;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:36, the fragment comprising eight contiguous amino acids of SEQ ID NO:36;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:35.

45. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:36;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:36, the fragment comprising eight contiguous amino acids of SEQ ID NO:36; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vc50\_1 deposited with the ATCC under accession number 98862;
- the protein being substantially free from other mammalian proteins.

46. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:37;
- (b) the nucleotide sequence of SEQ ID NO:37 from nucleotide 139 to nucleotide 1308;
- (c) the nucleotide sequence of SEQ ID NO:37 from nucleotide 211 to nucleotide 1308;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vc51\_1 deposited with the ATCC under accession number 98862;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vc51\_1 deposited with the ATCC under accession number 98862;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vc51\_1 deposited with the ATCC under accession number 98862;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vc51\_1 deposited with the ATCC under accession number 98862;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:38;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:38, the fragment comprising eight contiguous amino acids of SEQ ID NO:38;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and



(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:37.

47. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:38;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:38, the fragment comprising eight contiguous amino acids of SEQ ID NO:38; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vc51\_1 deposited with the ATCC under accession number 98862;
- the protein being substantially free from other mammalian proteins.

48. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:39;
- (b) the nucleotide sequence of SEQ ID NO:39 from nucleotide 21 to nucleotide 1142;
- (c) the nucleotide sequence of SEQ ID NO:39 from nucleotide 114 to nucleotide 1142;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vc52\_1 deposited with the ATCC under accession number 98862;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vc52\_1 deposited with the ATCC under accession number 98862;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vc52\_1 deposited with the ATCC under accession number 98862;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vc52\_1 deposited with the ATCC under accession number 98862;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:40;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:40, the fragment comprising eight contiguous amino acids of SEQ ID NO:40;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:39.

49. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:40;

(b) a fragment of the amino acid sequence of SEQ ID NO:40, the fragment comprising eight contiguous amino acids of SEQ ID NO:40; and

(c) the amino acid sequence encoded by the cDNA insert of clone vc52\_1 deposited with the ATCC under accession number 98862;

the protein being substantially free from other mammalian proteins.

50. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:41;

(b) the nucleotide sequence of SEQ ID NO:41 from nucleotide 13 to nucleotide 1416;

(c) the nucleotide sequence of SEQ ID NO:41 from nucleotide 346 to nucleotide 1416;

(d) the nucleotide sequence of the full-length protein coding sequence of clone vc33\_1 deposited with the ATCC under accession number 98886;

(e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vc33\_1 deposited with the ATCC under accession number 98886;

(f) the nucleotide sequence of a mature protein coding sequence of clone vc33\_1 deposited with the ATCC under accession number 98886;

(g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vc33\_1 deposited with the ATCC under accession number 98886;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:42;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:42, the fragment comprising eight contiguous amino acids of SEQ ID NO:42;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:41.

51. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:42;

(b) a fragment of the amino acid sequence of SEQ ID NO:42, the fragment comprising eight contiguous amino acids of SEQ ID NO:42; and

(c) the amino acid sequence encoded by the cDNA insert of clone vc33\_1 deposited with the ATCC under accession number 98886;

the protein being substantially free from other mammalian proteins.

52. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:43;

(b) the nucleotide sequence of SEQ ID NO:43 from nucleotide 232 to nucleotide 1461;

(c) the nucleotide sequence of SEQ ID NO:43 from nucleotide 280 to nucleotide 1461;

(d) the nucleotide sequence of the full-length protein coding sequence of clone vc34\_1 deposited with the ATCC under accession number 98886;

(e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vc34\_1 deposited with the ATCC under accession number 98886;

(f) the nucleotide sequence of a mature protein coding sequence of clone vc34\_1 deposited with the ATCC under accession number 98886;

(g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vc34\_1 deposited with the ATCC under accession number 98886;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:44;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:44, the fragment comprising eight contiguous amino acids of SEQ ID NO:44;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:43.

53. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:44;

(b) a fragment of the amino acid sequence of SEQ ID NO:44, the fragment comprising eight contiguous amino acids of SEQ ID NO:44; and

(c) the amino acid sequence encoded by the cDNA insert of clone vc34\_1 deposited with the ATCC under accession number 98886;

the protein being substantially free from other mammalian proteins.

54. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:45;

- (b) the nucleotide sequence of SEQ ID NO:45 from nucleotide 1922 to nucleotide 2350;
- (c) the nucleotide sequence of SEQ ID NO:45 from nucleotide 2237 to nucleotide 2350;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vc47\_1 deposited with the ATCC under accession number 98886;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vc47\_1 deposited with the ATCC under accession number 98886;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vc47\_1 deposited with the ATCC under accession number 98886;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vc47\_1 deposited with the ATCC under accession number 98886;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:46;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:46, the fragment comprising eight contiguous amino acids of SEQ ID NO:46;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:45.

55. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:46;
- (b) a fragment of the amino acid sequence of SEQ ID NO:46, the fragment comprising eight contiguous amino acids of SEQ ID NO:46; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vc47\_1 deposited with the ATCC under accession number 98886;

the protein being substantially free from other mammalian proteins.

56. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:47;
- (b) the nucleotide sequence of SEQ ID NO:47 from nucleotide 111 to nucleotide 1337;
- (c) the nucleotide sequence of SEQ ID NO:47 from nucleotide 246 to nucleotide 1337;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vc54\_1 deposited with the ATCC under accession number 98886;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vc54\_1 deposited with the ATCC under accession number 98886;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vc54\_1 deposited with the ATCC under accession number 98886;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vc54\_1 deposited with the ATCC under accession number 98886;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:48;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:48, the fragment comprising eight contiguous amino acids of SEQ ID NO:48;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:47.

57. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:48;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:48, the fragment comprising eight contiguous amino acids of SEQ ID NO:48; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vc54\_1 deposited with the ATCC under accession number 98886;
- the protein being substantially free from other mammalian proteins.

58. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:49;
- (b) the nucleotide sequence of SEQ ID NO:49 from nucleotide 189 to nucleotide 1637;
- (c) the nucleotide sequence of SEQ ID NO:49 from nucleotide 270 to nucleotide 1637;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vc57\_1 deposited with the ATCC under accession number 98886;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vc57\_1 deposited with the ATCC under accession number 98886;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vc57\_1 deposited with the ATCC under accession number 98886;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vc57\_1 deposited with the ATCC under accession number 98886;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:50;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:50; the fragment comprising eight contiguous amino acids of SEQ ID NO:50;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees

C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:49.

59. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:50;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:50, the fragment comprising eight contiguous amino acids of SEQ ID NO:50; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vc57\_1 deposited with the ATCC under accession number 98886;
- the protein being substantially free from other mammalian proteins.

60. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:51;
- (b) the nucleotide sequence of SEQ ID NO:51 from nucleotide 15 to nucleotide 1934;
- (c) the nucleotide sequence of SEQ ID NO:51 from nucleotide 1704 to nucleotide 1934;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone ve13\_1 deposited with the ATCC under accession number 98886;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone ve13\_1 deposited with the ATCC under accession number 98886;
- (f) the nucleotide sequence of a mature protein coding sequence of clone ve13\_1 deposited with the ATCC under accession number 98886;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone ve13\_1 deposited with the ATCC under accession number 98886;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:52;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:52, the fragment comprising eight contiguous amino acids of SEQ ID NO:52;



(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:51.

61. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:52;

(b) a fragment of the amino acid sequence of SEQ ID NO:52, the fragment comprising eight contiguous amino acids of SEQ ID NO:52; and

(c) the amino acid sequence encoded by the cDNA insert of clone ve13\_1 deposited with the ATCC under accession number 98886;

the protein being substantially free from other mammalian proteins.

62. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:53;

(b) the nucleotide sequence of SEQ ID NO:53 from nucleotide 240 to nucleotide 503;

(c) the nucleotide sequence of SEQ ID NO:53 from nucleotide 318 to nucleotide 503;

(d) the nucleotide sequence of the full-length protein coding sequence of clone ve16\_1 deposited with the ATCC under accession number 98886;

(e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone ve16\_1 deposited with the ATCC under accession number 98886;

(f) the nucleotide sequence of a mature protein coding sequence of clone ve16\_1 deposited with the ATCC under accession number 98886;

(g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone ve16\_1 deposited with the ATCC under accession number 98886;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:54;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:54, the fragment comprising eight contiguous amino acids of SEQ ID NO:54;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:53.

63. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:54;

(b) a fragment of the amino acid sequence of SEQ ID NO:54, the fragment comprising eight contiguous amino acids of SEQ ID NO:54; and

(c) the amino acid sequence encoded by the cDNA insert of clone ve16\_1 deposited with the ATCC under accession number 98886;

the protein being substantially free from other mammalian proteins.

64. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:55;

(b) the nucleotide sequence of SEQ ID NO:55 from nucleotide 11 to nucleotide 1063;

(c) the nucleotide sequence of SEQ ID NO:55 from nucleotide 71 to nucleotide 1063;

(d) the nucleotide sequence of the full-length protein coding sequence of clone vf3\_1 deposited with the ATCC under accession number 98886;

(e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vf3\_1 deposited with the ATCC under accession number 98886;

- (f) the nucleotide sequence of a mature protein coding sequence of clone vf3\_1 deposited with the ATCC under accession number 98886;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vf3\_1 deposited with the ATCC under accession number 98886;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:56;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:56, the fragment comprising eight contiguous amino acids of SEQ ID NO:56;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:55.

65. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:56;
- (b) a fragment of the amino acid sequence of SEQ ID NO:56, the fragment comprising eight contiguous amino acids of SEQ ID NO:56; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vf3\_1 deposited with the ATCC under accession number 98886;

the protein being substantially free from other mammalian proteins.

66. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:57;
- (b) the nucleotide sequence of SEQ ID NO:57 from nucleotide 542 to nucleotide 886;
- (c) the nucleotide sequence of SEQ ID NO:57 from nucleotide 755 to nucleotide 886;

(d) the nucleotide sequence of the full-length protein coding sequence of clone vj2\_1 deposited with the ATCC under accession number 98886;

(e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vj2\_1 deposited with the ATCC under accession number 98886;

(f) the nucleotide sequence of a mature protein coding sequence of clone vj2\_1 deposited with the ATCC under accession number 98886;

(g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vj2\_1 deposited with the ATCC under accession number 98886;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:58;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:58, the fragment comprising eight contiguous amino acids of SEQ ID NO:58;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:57.

67. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:58;

(b) a fragment of the amino acid sequence of SEQ ID NO:58, the fragment comprising eight contiguous amino acids of SEQ ID NO:58; and

(c) the amino acid sequence encoded by the cDNA insert of clone vj2\_1 deposited with the ATCC under accession number 98886;

the protein being substantially free from other mammalian proteins.

68. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:59;
- (b) the nucleotide sequence of SEQ ID NO:59 from nucleotide 30 to nucleotide 344;
- (c) the nucleotide sequence of SEQ ID NO:59 from nucleotide 84 to nucleotide 344;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vp7\_1 deposited with the ATCC under accession number 98886;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vp7\_1 deposited with the ATCC under accession number 98886;
- (f) the nucleotide sequence of a mature protein-coding sequence of clone vp7\_1 deposited with the ATCC under accession number 98886;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vp7\_1 deposited with the ATCC under accession number 98886;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:60;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:60, the fragment comprising eight contiguous amino acids of SEQ ID NO:60;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:59.

69. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:60;
- (b) a fragment of the amino acid sequence of SEQ ID NO:60, the fragment comprising eight contiguous amino acids of SEQ ID NO:60; and

(c) the amino acid sequence encoded by the cDNA insert of clone vp7\_1 deposited with the ATCC under accession number 98886; the protein being substantially free from other mammalian proteins.

70. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:61;
- (b) the nucleotide sequence of SEQ ID NO:61 from nucleotide 23 to nucleotide 757;
- (c) the nucleotide sequence of SEQ ID NO:61 from nucleotide 119 to nucleotide 757;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vp8\_1 deposited with the ATCC under accession number 98886;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vp8\_1 deposited with the ATCC under accession number 98886;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vp8\_1 deposited with the ATCC under accession number 98886;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vp8\_1 deposited with the ATCC under accession number 98886;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:62;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:62, the fragment comprising eight contiguous amino acids of SEQ ID NO:62;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:61.

71. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:62;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:62, the fragment comprising eight contiguous amino acids of SEQ ID NO:62; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vp8\_1 deposited with the ATCC under accession number 98886;
- the protein being substantially free from other mammalian proteins.

72. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:63;
- (b) the nucleotide sequence of SEQ ID NO:63 from nucleotide 1048 to nucleotide 3726;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone vb22\_1 deposited with the ATCC under accession number 98933;
- (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vb22\_1 deposited with the ATCC under accession number 98933;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:64;
- (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:64, the fragment comprising eight contiguous amino acids of SEQ ID NO:64;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:63.

73. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:64;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:64, the fragment comprising eight contiguous amino acids of SEQ ID NO:64; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vb22\_1 deposited with the ATCC under accession number 98933;
- the protein being substantially free from other mammalian proteins.

74. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:65;
- (b) the nucleotide sequence of SEQ ID NO:65 from nucleotide 134 to nucleotide 667;
- (c) the nucleotide sequence of SEQ ID NO:65 from nucleotide 191 to nucleotide 667;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vc48\_1 deposited with the ATCC under accession number 98933;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vc48\_1 deposited with the ATCC under accession number 98933;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vc48\_1 deposited with the ATCC under accession number 98933;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vc48\_1 deposited with the ATCC under accession number 98933;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:66;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:66, the fragment comprising eight contiguous amino acids of SEQ ID NO:66;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees



C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:65.

75. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:66;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:66, the fragment comprising eight contiguous amino acids of SEQ ID NO:66; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vc48\_1 deposited with the ATCC under accession number 98933;
- the protein being substantially free from other mammalian proteins.

76. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:67;
- (b) the nucleotide sequence of SEQ ID NO:67 from nucleotide 65 to nucleotide 457;
- (c) the nucleotide sequence of SEQ ID NO:67 from nucleotide 158 to nucleotide 457;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vp3\_1 deposited with the ATCC under accession number 98933;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vp3\_1 deposited with the ATCC under accession number 98933;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vp3\_1 deposited with the ATCC under accession number 98933;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vp3\_1 deposited with the ATCC under accession number 98933;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:68;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:68, the fragment comprising eight contiguous amino acids of SEQ ID NO:68;

- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:67.

77. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:68;
  - (b) a fragment of the amino acid sequence of SEQ ID NO:68, the fragment comprising eight contiguous amino acids of SEQ ID NO:68; and
  - (c) the amino acid sequence encoded by the cDNA insert of clone vp3\_1 deposited with the ATCC under accession number 98933;
- the protein being substantially free from other mammalian proteins.

78. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:69;
- (b) the nucleotide sequence of SEQ ID NO:69 from nucleotide 29 to nucleotide 1387;
- (c) the nucleotide sequence of SEQ ID NO:69 from nucleotide 113 to nucleotide 1387;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vc61\_1 deposited with the ATCC under accession number 207012;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vc61\_1 deposited with the ATCC under accession number 207012;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vc61\_1 deposited with the ATCC under accession number 207012;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vc61\_1 deposited with the ATCC under accession number 207012;

(h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:70;

(i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:70, the fragment comprising eight contiguous amino acids of SEQ ID NO:70;

(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:69.

79. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:70;

(b) a fragment of the amino acid sequence of SEQ ID NO:70, the fragment comprising eight contiguous amino acids of SEQ ID NO:70; and

(c) the amino acid sequence encoded by the cDNA insert of clone vc61\_1 deposited with the ATCC under accession number 207012;

the protein being substantially free from other mammalian proteins.

80. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:71;

(b) the nucleotide sequence of SEQ ID NO:71 from nucleotide 44 to nucleotide 1513;

(c) the nucleotide sequence of SEQ ID NO:71 from nucleotide 92 to nucleotide 1513;

(d) the nucleotide sequence of SEQ ID NO:71 from nucleotide 1 to nucleotide 458;

(e) the nucleotide sequence of the full-length protein coding sequence of clone vp15\_1 deposited with the ATCC under accession number 207012;

(f) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vp15\_1 deposited with the ATCC under accession number 207012;

(g) the nucleotide sequence of a mature protein coding sequence of clone vp15\_1 deposited with the ATCC under accession number 207012;

(h) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vp15\_1 deposited with the ATCC under accession number 207012;

(i) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:72;

(j) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:72, the fragment comprising eight contiguous amino acids of SEQ ID NO:72;

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(h); and

(l) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(h), and that has a length that is at least 25% of the length of SEQ ID NO:71.

81. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:72;

(b) the amino acid sequence of SEQ ID NO:72 from amino acid 1 to amino acid 139;

(c) a fragment of the amino acid sequence of SEQ ID NO:72, the fragment comprising eight contiguous amino acids of SEQ ID NO:72; and

(d) the amino acid sequence encoded by the cDNA insert of clone vp15\_1 deposited with the ATCC under accession number 207012;

the protein being substantially free from other mammalian proteins.

82. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:73;
- (b) the nucleotide sequence of SEQ ID NO:73 from nucleotide 348 to nucleotide 743;
- (c) the nucleotide sequence of SEQ ID NO:73 from nucleotide 414 to nucleotide 743;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vp17\_1 deposited with the ATCC under accession number 207012;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vp17\_1 deposited with the ATCC under accession number 207012;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vp17\_1 deposited with the ATCC under accession number 207012;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vp17\_1 deposited with the ATCC under accession number 207012;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:74;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:74, the fragment comprising eight contiguous amino acids of SEQ ID NO:74;
- (j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:73.

83. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:74;
- (b) a fragment of the amino acid sequence of SEQ ID NO:74, the fragment comprising eight contiguous amino acids of SEQ ID NO:74; and

(c) the amino acid sequence encoded by the cDNA insert of clone vp17\_1 deposited with the ATCC under accession number 207012; the protein being substantially free from other mammalian proteins.

84. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:75;
- (b) the nucleotide sequence of SEQ ID NO:75 from nucleotide 144 to nucleotide 461;
- (c) the nucleotide sequence of the full-length protein coding sequence of clone vp19\_1 deposited with the ATCC under accession number 207012;
- (d) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vp19\_1 deposited with the ATCC under accession number 207012;
- (e) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:76;
- (f) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:76, the fragment comprising eight contiguous amino acids of SEQ ID NO:76;
- (g) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d); and
- (h) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(d), and that has a length that is at least 25% of the length of SEQ ID NO:75.

85. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:76;
- (b) a fragment of the amino acid sequence of SEQ ID NO:76, the fragment comprising eight contiguous amino acids of SEQ ID NO:76; and
- (c) the amino acid sequence encoded by the cDNA insert of clone vp19\_1 deposited with the ATCC under accession number 207012;

the protein being substantially free from other mammalian proteins.

86. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:77;
- (b) the nucleotide sequence of SEQ ID NO:77 from nucleotide 54 to nucleotide 368;
- (c) the nucleotide sequence of SEQ ID NO:77 from nucleotide 141 to nucleotide 368;
- (d) the nucleotide sequence of SEQ ID NO:77 from nucleotide 51 to nucleotide 332;
- (e) the nucleotide sequence of the full-length protein coding sequence of clone vq1\_1 deposited with the ATCC under accession number 207012;
- (f) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vq1\_1 deposited with the ATCC under accession number 207012;
- (g) the nucleotide sequence of a mature protein coding sequence of clone vq1\_1 deposited with the ATCC under accession number 207012;
- (h) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vq1\_1 deposited with the ATCC under accession number 207012;
- (i) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:78;
- (j) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:78, the fragment comprising eight contiguous amino acids of SEQ ID NO:78;
- (k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(h); and
- (l) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(h), and that has a length that is at least 25% of the length of SEQ ID NO:77.

87. A protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:78;
  - (b) the amino acid sequence of SEQ ID NO:78 from amino acid 1 to amino acid 93;
  - (c) a fragment of the amino acid sequence of SEQ ID NO:78, the fragment comprising eight contiguous amino acids of SEQ ID NO:78; and
  - (d) the amino acid sequence encoded by the cDNA insert of clone vq1\_1 deposited with the ATCC under accession number 207012;
- the protein being substantially free from other mammalian proteins.

88. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:79;
- (b) the nucleotide sequence of SEQ ID NO:79 from nucleotide 2 to nucleotide 1018;
- (c) the nucleotide sequence of SEQ ID NO:79 from nucleotide 53 to nucleotide 1018;
- (d) the nucleotide sequence of the full-length protein coding sequence of clone vp14\_1 deposited with the ATCC under accession number 207011;
- (e) a nucleotide sequence encoding the full-length protein encoded by the cDNA insert of clone vp14\_1 deposited with the ATCC under accession number 207011;
- (f) the nucleotide sequence of a mature protein coding sequence of clone vp14\_1 deposited with the ATCC under accession number 207011;
- (g) a nucleotide sequence encoding a mature protein encoded by the cDNA insert of clone vp14\_1 deposited with the ATCC under accession number 207011;
- (h) a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID NO:80;
- (i) a nucleotide sequence encoding a protein comprising a fragment of the amino acid sequence of SEQ ID NO:80, the fragment comprising eight contiguous amino acids of SEQ ID NO:80;



(j) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 65 degrees C, or 4X SSC at 42 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g); and

(k) the nucleotide sequence of a polynucleotide that hybridizes under conditions at least as stringent as 4X SSC at 50 degrees C, or 6X SSC at 40 degrees C with 50% formamide, to any one of the polynucleotides specified by (a)-(g), and that has a length that is at least 25% of the length of SEQ ID NO:79.

89. A protein comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:80;

(b) a fragment of the amino acid sequence of SEQ ID NO:80, the fragment comprising eight contiguous amino acids of SEQ ID NO:80; and

(c) the amino acid sequence encoded by the cDNA insert of clone vp14\_1 deposited with the ATCC under accession number 207011;

the protein being substantially free from other mammalian proteins.



Clone "vb21\_1"

A polynucleotide of the present invention has been identified as clone "vb21\_1". vb21\_1 was isolated from a human fetal brain cDNA library and was identified as encoding a secreted or transmembrane protein on the basis of computer analysis of the amino acid sequence of the encoded protein. vb21\_1 is a full-length clone, including the entire coding sequence of a secreted protein (also referred to herein as "vb21\_1 protein").

The nucleotide sequence of vb21\_1 as presently determined is reported in SEQ ID NO:19, and includes a poly(A) tail. What applicants presently believe to be the proper reading frame and the predicted amino acid sequence of the vb21\_1 protein corresponding to the foregoing nucleotide sequence is reported in SEQ ID NO:20. Amino acids 296 to 308 of SEQ ID NO:20 are a predicted leader/signal sequence, with the predicted mature amino acid sequence beginning at amino acid 309. Due to the hydrophobic nature of the predicted leader/signal sequence, it is likely to act as a transmembrane domain should the predicted leader/signal sequence not be separated from the remainder of the vb21\_1 protein.

The EcoRI/NotI restriction fragment obtainable from the deposit containing clone vb21\_1 should be approximately 4159 bp.

The nucleotide sequence disclosed herein for vb21\_1 was searched against the GenBank and GeneSeq nucleotide sequence databases using BLASTN/BLASTX and FASTA search protocols. vb21\_1 demonstrated at least some similarity with sequences identified as AA026150 (zj99c10.s1 Soares pregnant uterus NbHPU Homo sapiens cDNA clone 469170 3', mRNA sequence), T72108 (Human semaphorin Z gene; standard; cDNA to mRNA), U52840 (Human semaphorin F homolog), X97817 (M. musculus mRNA for semaphorin F), and X97818 (M. musculus mRNA for semaphorin G). The predicted amino acid sequence disclosed herein for vb21\_1 was searched against the GenPept and GeneSeq amino acid sequence databases using the BLASTX search protocol. The predicted vb21\_1 protein demonstrated at least some similarity to sequences identified as W19857 (Human semaphorin Z) and X97818 (samaphorin G [Mus musculus]). Semaphorins are important membrane proteins involved in axonal guidance in the embryonic stage, and may also have a role in nerve regeneration after injury. Based upon sequence similarity, vb21\_1 proteins and each similar protein or peptide may share at least some activity. The TopPredII computer program predicts four additional potential transmembrane domains within the vb21\_1 protein sequence, centered around amino acids 237, 523, 769, and 895 of SEQ ID NO:20, respectively.



&lt;400&gt; 18

Met Glu Leu Pro Ala Val Asn Leu Lys Val Ile Leu Leu Gly His Trp  
 1 5 10 15

Leu Leu Thr Thr Trp Gly Cys Ile Val Phe Ser Gly Ser Tyr Ala Trp  
 20 25 30

Ala Asn Phe Thr Ile Leu Ala Leu Gly Val Trp Ala Val Ala Gln Arg  
 35 40 45

Asp Ser Ile Asp Ala Ile Ser Met Phe Leu Gly Gly Leu Leu Ala Thr  
 50 55 60

Ile Phe Leu Asp Ile Val His Ile Ser Ile Phe Tyr Pro Arg Val Ser  
 65 70 75 80

Leu Thr Asp Thr Gly Arg Phe Gly Val Gly Met Ala Ile Leu Ser Leu  
 85 90 95

Leu Leu Lys Pro Leu Ser Cys Cys Phe Val Tyr His Met Tyr Arg Glu  
 100 105 110

Arg Gly Gly Phe Leu Gly Ser Ser Gln Asp Arg Ser Ala Tyr Gln Thr  
 115 120 125

Ile Asp Ser Ala Glu Ala Pro Ala Asp Pro Phe Ala Val Pro Glu Gly  
 130 135 140

Arg Ser Gln Asp Ala Arg Gly Tyr  
 145 150

&lt;210&gt; 19

&lt;211&gt; 4144

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 19

ctccttgggc aggtgaggat aacagccctg tggcgagctg gtcctgagca tctgagttag 60  
 cagtgatgcg gagaaatgac cttgggtggga ggggacgtcc acagacgtga cccgggacgtc 120  
 gggacagggtg gatctgggggt caagggggagt gtttagaata cttggtggca tccatgacgc 180  
 agtgggtgtga gcgtgagata tccagcatcg cacctggaga gctgtgctgt ctgctccttt 240  
 ccttcctgcc tcaggaggag tgtcagaact acgtgagagt cctgatcgtc gccggccgga 300  
 aggtgttcat gtgtggaacc aatgcctttt ccccatgtg caccagcaga caggtgggga 360  
 acctcagccg gactattgag aagatcaatg gtgtggcccg ctgcccctat gaccacgccc 420  
 acaactccac agctgtcatc tcctcccagg gggagctcta tgcagccacg gtcacgact 480  
 tctcaggtcg ggaccctgcc atctaccgca gcctgggcag tgggcccaccg ctccgactcg 540  
 cccaatataa ctccaagtgg cttaatgagc caaacttcgt ggcagcctat gatattgggc 600  
 tgtttgcata cttcttctctg cgggagaacg cagtggagca cgactgtgga cgcaccgtgt 660  
 actctcgcgt ggcccgctg tgcagaatg acgtgggggg ccgattcctg ctggaggaca 720  
 catggaccac attcatgaag gcccggtcga actgctccc cccgggagag gtccccttct 780  
 actataacga gctgcagagt gccttccact tgccggagca ggacctcatc tatggagttt 840  
 tcacaaccaa cgtaaacagc attgcggtt ctgctgtctg cgccttcaac ctcaagtcta 900  
 tctcccaggc tttcaatggc ccatttcgct accaggagaa ccccagggtt gcctggctcc 960  
 ccatagccaa ccccatcccc aatttccagt gtggcaccct gcctgagacc ggtcccaacg 1020  
 agaacctgac ggagcgcagc ctgcaggagc cgcagcgcct ctccctgatg agcgaggccg 1080  
 tgcagccggt gacacccgag ccctgtgtca cccaggacag cgtgcgcttc tcacacctcg 1140  
 tgggtggacct ggtgcaggct aaagacacgc tctaccatgt actctacatt ggcaccgagt 1200  
 cgggcaccat cctgaaggcg ctgtccacgg cgagccgcag cctccacggc tgctacctgg 1260



```

aggagctgca cgtgctgccc cccggggcgc ggcagcccct gcgcagcctg cgcattcctgc 1320
acagcgcccg cgcgctcttc gtggggctga gagacggcgt cctgcggggc ccactggaga 1380
ggtgcgccgc ctaccgcagc cagggggcat gcctgggggc cggggacccg tactgtggct 1440
gggacgggaa gcagcaacgt tgcagcacac tcgaggacag ctccaacatg agcctctgga 1500
cccagaacat caccgcctgt cctgtgcgga atgtgacacg ggatgggggc ttcgggccc 1560
ggtcaccatg gcaaccatgt gagcacttgg atggggacaa ctcaggctct tgcctgtgtc 1620
gagctcgatc ctgtgattcc cctcgacccc gctgtggggg ccttgactgc ctggggccag 1680
ccatccacat cgccaactgc tccaggaatg gggcgtggac cccgtggtca tcgtgggcgc 1740
tgtgcagcac gtccctgtggc atcggttcc aggtccgcca gcgaagtgc agcaacctgc 1800
ctccccgcca cggggggcgc atctgctgg gcaagagcgc ggaggaaagg ttctgtaatg 1860
agaacacgcc ttgcccgggt cccatcttct gggcttccct gggctcctgg agcaagtgc 1920
gcagcaactg tggagggggc atgcagtcgc ggcgtcgggc ctgcgagaa ggcaactcct 1980
gcttgggctg cggcgtggag tcaagacgt gcaacccgga gggctgcccc gaagtgcggc 2040
gcaacacccc ctggacgccc tggctgcccc tgaacgtgac gcagggcggg gcacggcagg 2100
agcagcggtt ccgcttcacc tgcgcgcgc ccttgcaga cccgcacggc ctgcagttcg 2160
gcaggagaag gaccgagacg aggcctgtc ccgcggacgg ctccggctcc tgcgacacg 2220
acgtggcctg ggaggacctc ctgcgcagcg ggagcaacct cccgcacacg gtgagcggg 2280
gctggggcgc ctggggccc tggctgctct gctcccgga ctgcgagctg ggcttccgcg 2340
tccgcgaagag aactgactc aacccggag cccgcaacgg gggcctgccc tgcgtgggcg 2400
atgttgccga gtaccaggac tgcaaccccc aggttgcgc agttcggggg gcttggctct 2460
ggtgcacctc atggtctcca tgcagctt cctgtgggtg gggctactat caacgcacc 2520
gttctgcac cagccccgca cctccccag gtgaggacat ctgtctcggg ctgcacacg 2580
aggaggcact atgtgccaca caggcctgcc cagaaggctg gtcqccctgg tctgagtga 2640
gtaagtgcac tgacgacgga gccagagcgc gaagccggca ctgtgaggag ctctccccag 2700
gggtccagcg cgtgtctgga aacagcagcc agagccgccc ctgcccctac agcgagattc 2760
ccgtcactct gccagcctc agcatggagg aggcacccg ctgtgcaggg ttcaatctca 2820
tccacttggt ggccacgggc atctcctgct tcttgggctc tgggctcctg acctagcag 2880
tgtacctgtc ttgcccagc tgcagcgtc agtccaggga gtccacactg gtccatcctg 2940
ccacccccaa ccatttgcac tacaagggcg gaggcacccc gaagaatgaa aagtacacac 3000
ccatggaatt caagaccctg aacaagaata acttgatccc tgatgacaga gccacttct 3060
accatttga gcagaccaat gtgtacacga ctacttacta cccaagcccc ctgaacaaac 3120
acagcttcgc gccgaggcc tcacctggac aacggtgctt cccaacagc tgataccgc 3180
gtcctgggga cttgggcttc ttgccttcat aaggcacaga gcagatggag atgggacagt 3240
ggagccagtt tggttttct cctctgact aggcacagaa cttgctgcct tgcctgtgg 3300
gggtccctac cggcttcaga gagctctggc tggcattgac catgggggaa agggctggt 3360
tcaggctgac atatggcgc aggtccagtt cagcccaggt ctctcatggt tatcttcaa 3420
cccactgtca cgctgacact atgtgccc atgctgggct tggacctact gggcatttga 3480
gggaattggag aatggagatg gcaagagggc aggtctttaa gtttgggtg gagacaact 3540
cctgtggccc ccacaagctg agtctggcct tctccagctg gccccaaaaa aggcctttgc 3600
tacatcctga ttatctctga aagtaatcaa tcaagtggct ccagtagctc tggattttct 3660
gccagggctg ggccattgtg gtgctgcccc agtatgacat gggaccaagg ccagcgagg 3720
ttatccacct ctgcctggaa gtctatactc taccagggc atccctctgg tcagaggcag 3780
tgagtactgg gaactggagg ctgacctgtg cttagaagtc ctttaactct ggctgggtaca 3840
ggcctcagcc ttgcccctaa tgcacgaaag gtggccagg agagaggatc aatgccatag 3900
gagggcagaag tctggcctct gtgcctctat ggagactatc ttccagttgc tgctcaacag 3960
agttgttggc tgagacctgc ttgggagttc ctgctggccc ttcattctgt caggaaacaca 4020
cacacacaca cactcacaca cgcacacaca atcacattt gctacagcaa caaaaaagac 4080
attgggctgt ggcattatta attaaagatg atatccagtc aaaaaaaaaa aaaaaaaaaa 4144
aaaa

```

&lt;210&gt; 20

&lt;211&gt; 999

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 20

Met Thr Gln Trp Cys Glu Arg Glu Ile Ser Ser Ile Ala Pro Gly Glu  
1 5 10 15

Leu Cys Cys Leu Leu Leu Ser Phe Leu Pro Gln Glu Glu Cys Gln Asn





			20						25								30		
Tyr	Val	Arg	Val	Leu	Ile	Val	Ala	Gly	Arg	Lys	Val	Phe	Met	Cys	Gly				
		35					40					45							
Thr	Asn	Ala	Phe	Ser	Pro	Met	Cys	Thr	Ser	Arg	Gln	Val	Gly	Asn	Leu				
	50					55					60								
Ser	Arg	Thr	Ile	Glu	Lys	Ile	Asn	Gly	Val	Ala	Arg	Cys	Pro	Tyr	Asp				
	65				70					75					80				
Pro	Arg	His	Asn	Ser	Thr	Ala	Val	Ile	Ser	Ser	Gln	Gly	Glu	Leu	Tyr				
				85					90					95					
Ala	Ala	Thr	Val	Ile	Asp	Phe	Ser	Gly	Arg	Asp	Pro	Ala	Ile	Tyr	Arg				
			100					105						110					
Ser	Leu	Gly	Ser	Gly	Pro	Pro	Leu	Arg	Thr	Ala	Gln	Tyr	Asn	Ser	Lys				
		115					120					125							
Trp	Leu	Asn	Glu	Pro	Asn	Phe	Val	Ala	Ala	Tyr	Asp	Ile	Gly	Leu	Phe				
	130					135					140								
Ala	Tyr	Phe	Phe	Leu	Arg	Glu	Asn	Ala	Val	Glu	His	Asp	Cys	Gly	Arg				
	145				150					155					160				
Thr	Val	Tyr	Ser	Arg	Val	Ala	Arg	Val	Cys	Lys	Asn	Asp	Val	Gly	Gly				
				165					170					175					
Arg	Phe	Leu	Leu	Glu	Asp	Thr	Trp	Thr	Thr	Phe	Met	Lys	Ala	Arg	Leu				
			180					185					190						
Asn	Cys	Ser	Arg	Pro	Gly	Glu	Val	Pro	Phe	Tyr	Tyr	Asn	Glu	Leu	Gln				
		195					200					205							
Ser	Ala	Phe	His	Leu	Pro	Glu	Gln	Asp	Leu	Ile	Tyr	Gly	Val	Phe	Thr				
	210					215					220								
Thr	Asn	Val	Asn	Ser	Ile	Ala	Ala	Ser	Ala	Val	Cys	Ala	Phe	Asn	Leu				
	225				230					235					240				
Ser	Ala	Ile	Ser	Gln	Ala	Phe	Asn	Gly	Pro	Phe	Arg	Tyr	Gln	Glu	Asn				
				245				250						255					
Pro	Arg	Ala	Ala	Trp	Leu	Pro	Ile	Ala	Asn	Pro	Ile	Pro	Asn	Phe	Gln				
		260						265						270					
Cys	Gly	Thr	Leu	Pro	Glu	Thr	Gly	Pro	Asn	Glu	Asn	Leu	Thr	Glu	Arg				
		275					280					285							
Ser	Leu	Gln	Asp	Ala	Gln	Arg	Leu	Phe	Leu	Met	Ser	Glu	Ala	Val	Gln				
	290					295					300								
Pro	Val	Thr	Pro	Glu	Pro	Cys	Val	Thr	Gln	Asp	Ser	Val	Arg	Phe	Ser				
	305				310					315</									





980

985

990

Gln Arg Cys Phe Pro Asn Ser  
995

<210> 21  
<211> 2820  
<212> DNA  
<213> Homo sapiens

<400> 21  
agaatgggag ctgtcagtta tcattcagga caaaggcaat cctcagctac ataccaaaagt 60  
ccttctgaag tgcattgatct ttgaatatgc agagtcgggtg acaagtacag caatgacttc 120  
agtaagccag gcatccttgg atgtctccat gataataatt atttcccttag gagcaatttg 180  
tgcagtggtg ctggttatta tgggtctatt tgcaactagg tgtaaccgcg agaagaaaga 240  
cactagatcc tataactgca ggggtggccga atcaacttac cagcaccacc caaaaaggcc 300  
atcccgagc attcacaaag gggacatcac attggtgcct accataaatg gcactctgcc 360  
catcagatct catcacagat cgtctccatc ttcattctcct accttagaaa gagggcagat 420  
gggcagccgg cagagtcaca acagtcacca gtcactcaac agtttgggtga caatctcacc 480  
aaaccacgtg ccagagaatt tctcattaga actcaccac gccactcctg ctggttaggt 540  
ctctcagctt ctttcaatgc ttcaccagg gcaatatcag ccaagacca gttttcgagg 600  
aaacaaatat tccaggagct acagatatgc ctttcaagac atggacaaat ttagcttgaa 660  
agacagtggc cgtggtgaca gtgaggcagg agacagtgat tatgatttgg ggcgagattc 720  
tccaatagat aggtctgttg gtgaaggatt cagcgacctg tttctcacag atggaagaat 780  
tcagcagct atgagactct gcacggagg gtgcagggtc ctgggacact ctgaccagt 840  
ctggatgcca ccactgccct caccgtcttc tgattatagg agtaacatgt tcattccagg 900  
ggaagaattc ccaacgcaac ccagcagca gcattccacat cagagtcttg aggatgacgc 960  
tcagcctgca gattccgggtg aaaagaagaa gagtttttcc acctttggaa aggactcccc 1020  
aaacgatgag gacactgggg ataccagcac atcatctctg ctctcggaaa tgagcagtg 1080  
gttccagcgt ctcttaccgc cttccctgga cactattct gaatgcagtg aggtggatcg 1140  
gtccaaactc ctggagcgca ggaagggacc cttgccagcc aaactgtgg gttaccaca 1200  
gggggtagcg gcatgggcag ccagtcacga ttttcaaaat cccaccacca actgtgggcc 1260  
gccacttgga actcactcca gtgtgcagcc ttcttcaaaa tgggtgccag ccatggaggga 1320  
gattccctgaa aattatgagg aagatgattt tgacaatgtg ctcaaccacc tcaatgatgg 1380  
gaaacacgaa ctcatggatg ccagtgaact ggtggcagag attaacaac tgcttcaaga 1440  
tgtccgcccag agctaggaga ttttagcgaa gcatttttgt ttccatgtat atggaatatg 1500  
ggaacaacaa caacaacaaa aaaccctgaa agaactggca ttgccaaata gttgcattta 1560  
tcataaatgt gtctgtgtat attgaatatt aaatactgta ttttcgtatg tacacaatgc 1620  
aagtgtgatt attttaatct gtattttaaa aatacatttg taccttata tttatgtgta 1680  
ttaacaaac aaattttatt tttttactcc catgacagac atgtttttcc tagtctgtga 1740  
gaaactagcc actgttcaaa tctgatacac tattcaacca caaagtgtaa aggcactgct 1800  
tagattagtt ttgttgggga agaattatta tgttgatga acaaccccac tgaagcatta 1860  
tacaattctt aattccatta agtgatocca ctttttttca ataactttt agaaattaag 1920  
aatcattaaa attgttaagc tattttattg ttattttctc tactttctac tagccccaat 1980  
agttgaaactc ttataggaaa atcgaaagat aaagtgaag tttatttcag gactgagaaa 2040  
tatcttgaag gttattttatt agatgactat ctcaaatgaa ctttttatag acaatgatga 2100  
aaacagaaatt aaagtcaatg ttctctgact ccaggcccc tactattcca ggccatcaca 2160  
ctggcctgtt ccggagaata tttctctcac aatattatta tctacttata attatggtaa 2220  
acaataaaatt ttattccatc cttgtagtat gaaacatgct ccaaggaaat ggaatctgtc 2280  
ctttaaattg ataacagtat gtgttctaatt ggcataaaat attactggat aaaaacagtt 2340  
gtgtcagtg etctcctaag gtatgaaata taattgactt attctgaacc cattctattt 2400  
tgaatctccc ctttctctc acaatacttg aacattttta tcttttggaa tattgtcttt 2460  
ctttgttata actattcatt tttagctttt gtctccagt catgatctca tatttttgc 2520  
tttattttta gtataagaac atttataaaa tcatattttt gttactgcaa ttgttttatt 2580  
tggtgtgtgg caaatgagaa atcctttatt tattgtgctg tgatctctct gtgtggaatg 2640  
cctgggtgag agagatgctt attatgacta ttatcatttc tgaccaagct tctattaatg 2700  
ttattttctaa taatacacta tcttgattgt actctccaga aaatttttct gtcagtga 2760  
ataaaagaaa aattaaagta aagctaagga actgtcaaaa aaaaaaaaaa aaaaaaaaaa 2820



660	665	670
Pro Ala Asp Gly Ser Gly Ser Cys Asp Thr Asp Ala Leu Val Glu Asp 675 680 685		
Leu Leu Arg Ser Gly Ser Thr Ser Pro His Thr Val Ser Gly Gly Trp 690 695 700		
Ala Ala Trp Gly Pro Trp Ser Ser Cys Ser Arg Asp Cys Glu Leu Gly 705 710 715 720		
Phe Arg Val Arg Lys Arg Thr Cys Thr Asn Pro Glu Pro Arg Asn Gly 725 730 735		
Gly Leu Pro Cys Val Gly Asp Ala Ala Glu Tyr Gln Asp Cys Asn Pro 740 745 750		
Gln Ala Cys Pro Val Arg Gly Ala Trp Ser Cys Trp Thr Ser Trp Ser 755 760 765		
Pro Cys Ser Ala Ser Cys Gly Gly Gly His Tyr Gln Arg Thr Arg Ser 770 775 780		
Cys Thr Ser Pro Ala Pro Ser Pro Gly Glu Asp Ile Cys Leu Gly Leu 785 790 795 800		
His Thr Glu Glu Ala Leu Cys Ala Thr Gln Ala Cys Pro Glu Gly Trp 805 810 815		
Ser Pro Trp Ser Glu Trp Ser Lys Cys Thr Asp Asp Gly Ala Gln Ser 820 825 830		
Arg Ser Arg His Cys Glu Glu Leu Leu Pro Gly Ser Ser Ala Cys Ala 835 840 845		
Gly Asn Ser Ser Gln Ser Arg Pro Cys Pro Tyr Ser Glu Ile Pro Val 850 855 860		
Ile Leu Pro Ala Ser Ser Met Glu Glu Ala Thr Gly Cys Ala Gly Phe 865 870 875 880		
Asn Leu Ile His Leu Val Ala Thr Gly Ile Ser Cys Phe Leu Gly Ser 885 890 895		
Gly Leu Leu Thr Leu Ala Val Tyr Leu Ser Cys Gln His Cys Gln Arg 900 905 910		
Gln Ser Gln Glu Ser Thr Leu Val His Pro Ala Thr Pro Asn His Leu 915 920 925		
His Tyr Lys Gly Gly Gly Thr Pro Lys Asn Glu Lys Tyr Thr Pro Met 930 935 940		
Glu Phe Lys Thr Leu Asn Lys Asn Asn Leu Ile Pro Asp Asp Arg Ala 945 950 955 960		
Asn Phe Tyr Pro Leu Gln Gln Thr Asn Val Tyr Thr Thr Tyr Tyr 965 970 975		
Pro Ser Pro Leu Asn Lys His Ser Phe Arg Pro Glu Ala Ser Pro Gly		



340					345					350					
Ala	Ser	Arg	Ser	Leu	His	Gly	Cys	Tyr	Leu	Glu	Glu	Leu	His	Val	Leu
	355						360					365			
Pro	Pro	Gly	Arg	Arg	Glu	Pro	Leu	Arg	Ser	Leu	Arg	Ile	Leu	His	Ser
	370					375					380				
Ala	Arg	Ala	Leu	Phe	Val	Gly	Leu	Arg	Asp	Gly	Val	Leu	Arg	Val	Pro
385					390					395					400
Leu	Glu	Arg	Cys	Ala	Ala	Tyr	Arg	Ser	Gln	Gly	Ala	Cys	Leu	Gly	Ala
				405					410					415	
Arg	Asp	Pro	Tyr	Cys	Gly	Trp	Asp	Gly	Lys	Gln	Gln	Arg	Cys	Ser	Thr
			420					425					430		
Leu	Glu	Asp	Ser	Ser	Asn	Met	Ser	Leu	Trp	Thr	Gln	Asn	Ile	Thr	Ala
		435					440					445			
Cys	Pro	Val	Arg	Asn	Val	Thr	Arg	Asp	Gly	Gly	Phe	Gly	Pro	Trp	Ser
450						455					460				
Pro	Trp	Gln	Pro	Cys	Glu	His	Leu	Asp	Gly	Asp	Asn	Ser	Gly	Ser	Cys
465					470					475					480
Leu	Cys	Arg	Ala	Arg	Ser	Cys	Asp	Ser	Pro	Arg	Pro	Arg	Cys	Gly	Gly
				485					490					495	
Leu	Asp	Cys	Leu	Gly	Pro	Ala	Ile	His	Ile	Ala	Asn	Cys	Ser	Arg	Asn
			500				505						510		
Gly	Ala	Trp	Thr	Pro	Trp	Ser	Ser	Trp	Ala	Leu	Cys	Ser	Thr	Ser	Cys
		515					520						525		
Gly	Ile	Gly	Phe	Gln	Val	Arg	Gln	Arg	Ser	Cys	Ser	Asn	Pro	Ala	Pro
530						535					540				
Arg	His	Gly	Gly	Arg	Ile	Cys	Val	Gly	Lys	Ser	Arg	Glu	Glu	Arg	Phe
545					550					555					560
Cys	Asn	Glu	Asn	Thr	Pro	Cys	Pro	Val	Pro	Ile	Phe	Trp	Ala	Ser	Trp
				565					570					575	
Gly	Ser	Trp	Ser	Lys	Cys	Ser	Ser	Asn	Cys	Gly	Gly	Gly	Met	Gln	Ser
			580				585						590		
Arg	Arg	Arg	Ala	Cys	Glu	Asn	Gly	Asn	Ser	Cys	Leu	Gly	Cys	Gly	Val
		595					600					605			
Glu	Phe	Lys	Thr	Cys	Asn	Pro	Glu	Gly	Cys	Pro	Glu	Val	Arg	Arg	Asn
610						615					620				
Thr	Pro	Trp	Thr	Pro	Trp	Leu	Pro	Val	Asn	Val	Thr	Gln	Gly	Gly	Ala
625					630					635					640
Arg	Gln	Glu	Gln	Arg	Phe	Arg	Phe	Thr	Cys	Arg	Ala	Pro	Leu	Ala	Asp
				645					650					655	
Pro	His	Gly	Leu	Gln	Phe	Gly	Arg	Arg	Arg	Thr	Glu	Thr	Arg	Thr	Cys